

**Mass-spectral Studies of the Chemical Nature of the Lachrymatory Factor  
Formed Enzymically from S-(1-propenyl)-cysteine Sulfoxide Isolated  
from Onion (*Allium cepa*)**

T. MOISIO, C.-G. SPÅRE, and ARTTURI I. VIRTANEN

*Laboratory of the Foundation for Chemical Research, Biochemical Institute, Helsinki, Finland*

Mass-spectral studies have given some evidence that propenylsulfenic acid is the lachrymatory factor in onion<sup>1</sup>. When a dialyzed enzyme preparation from onion was added to a dilute solution of S-(1-propenyl)-cysteine sulfoxide, a strong peak of mass 90 was found already after a reaction time of 1 min. After a reaction time of 1 h, the peak was already lower and the peak of mass 58 corresponding to propionaldehyde had become stronger. After 2 h the peak of mass 90 had almost disappeared.

A peak of mass 98 was also found after the reaction time of 1 h. 2-Methyl-2-pentenal (mol.wt. 98) is believed to cause this peak. This unsaturated aldehyde was earlier isolated in this laboratory as a volatile product of an enzymic reaction in crushed onion and was taken to be derived from propionaldehyde<sup>2,3</sup>.

According to the results obtained so far, the lachrymatory factor has a short life in water solution at room temperature. If the lachrymatory factor is really propenylsulfenic acid, this would be the first aliphatic sulfenic acid whose existence has been proved.

A more detailed paper will be published in *Acta Chemica Scandinavica*.

This investigation is a part of a research project under U. S. Public Law No. 480, 83rd Congress.

**References:**

1. Virtanen, A. I. and Spåre, C.-G. *Suomen Kemistilehti B* **35** (1962) 28.
2. Virtanen, A. I. and Spåre, C.-G. *Ibid.* **34** (1961) 18.
3. Spåre, C.-G. and Virtanen, A. I. *Acta Chem. Scand.* **15** (1961) 1280.